

Max Arocha, DMD
1125 S.University Dr. Plantation, FL 33324
Emergency number (954) 432- 7771

All-on-four for teeth-in-a-day Home Care Instructions

We would like to thank you for choosing us to perform your dental implant surgical procedure. If you have any questions or concerns, please contact our office or Dr. Arocha's mobile number provided to you on the day of the all-on-four dental implants treatment.

General Instructions

Occasionally, the teeth-in-a-day dental prosthesis cover screw or healing abutment may loosen or fall out. This should not affect the stability of your implant. It is important, however, to call us as soon as possible so we can replace it for you.

Pain Some degree of discomfort and pain arises as the numbness subsides. We recommend that you start taking a non-steroidal anti-inflammatory once you arrive home. Ibuprofen/Motrin are such medications and should be taken as directed on the bottle (usually 600 mg every 6-8 hours). If you are unable to take anti-inflammatory medications or the pain is not relieved, then you may take the prescribed pain medication as directed. All pain medications have the ability to cause severe nausea and vomiting, so try to have some food in your stomach before you take them.

Do not wait for the pain to become unbearable before using some form of pain medication, as then it will be more difficult to control. Moderate to severe pain from all-on-four dental implants treatment usually does not last longer than 24-72 hours, sometimes peaking on the 3rd postoperative day. Relief should begin on the 4th post-operative day.

Bleeding Expect minor bleeding or OOZING from the operative site. This bleeding may continue throughout the first day. Once the oozing has slowed down, try to avoid placing gauze or any other irritant near the surgical site as this will only lead to further bleeding and discomfort.

If active bleeding should recur at any time, carefully rinse your mouth with ice cold water. Should active bleeding increases or persist, please call the office and a member of our surgical team will return your call promptly.

The Surgical Site The first stages of healing are aided by allowing tissues to rest. Avoid vigorous chewing, excessive spitting, or rinsing for the first 24 hours as initial healing may be delayed, active bleeding restarted, or infection introduced.

Try to avoid smoking completely, as it tends to slow the healing process and may also contribute to infection & prolonged discomfort.

Do not worry about the stitches if they start to come out. The sutures usually dissolve in 5-7 days and typically do not require removal.

Swelling Swelling usually develops during the first 12-24 hours following surgery, often peaking on the 3rd post-operative day. Swelling can be minimized by placing an ice pack on the affected side at 30 minute intervals during the first 48 hours after All-on-four dental implant treatment.

Any swelling, soreness, or stiffness in the jaw muscles can be relieved by applying a warm moist towel to the affected side of the face several times a day. Moist heat should be used after the first 48 hours.

Activity Limit activity during the first 24-48 hours after dental implant surgery. Excessive exertion may lead to postoperative bleeding and discomfort. When you lie down, keep your head elevated at least 45 degrees on a pillow.

Do not drive an automobile for 24 hours following dental implant surgery if you have had general anesthesia or if you are taking prescription pain medication.

Diet FLUID INTAKE IS VERY IMPORTANT. You can advance to soft foods and other fluids as tolerated. Please avoid hot liquids until the numbness has worn off, and the bleeding has stopped.

Food Selection is largely a matter of your choice. Soft, cool foods that require little or no chewing are most easily tolerated.

Oral Hygiene Start cleaning your mouth 24 hours after surgery by using warm salt water rinses (1/4 teaspoon table salt with a small cup of water). Repeat 3 times daily for one week. BRUSH your teeth with a SOFT TOOTH brush starting on the first postoperative day being careful not to irritate the dental implant surgical site.

Your Medications Take any prescribed medication on your regular schedule unless advised to do otherwise by your physician. Patients taking blood-thinners (ie. Coumadin, Plavix) should restart those medications after surgery unless indicated otherwise by your physician.

IF YOU FEEL THAT HEALING IS NOT PROGRESSING IN A NORMAL FASHION CONTACT OUR OFFICE
AT (954-432-7771) A MEMBER OF OUR SURGICAL TEAM WILL RETURN YOUR CALL IN A TIMELY
FASHION.

IF YOU ARE EXPERIENCING A TRUE MEDICAL EMERGENCY, CALL 911 IMMEDIATELY.